

Vegetation Atmosphere INteractions (VAINT)

Evgenii Churiulin¹, Merja Tölle¹,
Juergen Helmert² and Jean-Marie Bettems³

¹ – Center for Environmental Systems Research, Kassel, Germany

² – Deutscher Wetterdienst, Offenbach am Main, Germany

³ – MeteoSwiss, Zurich, Switzerland

Motivation

Changes in the seasonal phenological cycle due to winter of

1) summer crops affect the local climate by changes in the biogeophysical processes;

Seasonal phenology of vegetation impacts on the energy and

2) water cycle, and can amplify extreme events by changes of the seasonal cycle of the albedo and water availability;

The frequency of extreme events increases and will increase in

3) the near future. Hence, the need for modelling phenology will also increase;

Motivation

Forecasting the phenological cycle of vegetation provides


- 4) valuable information to assist with turbulent flux estimations relevant for boundary layer processes;

Knowing the evolution and magnitude of the seasonal LAI and

- 5) the plant coverage allows for a more realistic estimation of the surface albedo;

The need to improve


*Example of annual cycle of leaf area index for **grass** in the COSMO model*



Jan-Peter Schulz et.al., 2015

The need to improve

*Example of annual cycle of leaf area index for **grass** in the COSMO model*



Jan-Peter Schulz et.al., 2015

Relevance

COSMO model use simplified phenology scheme, which are in

1) general not capable to model the complex processes of the growing season in spring, the evolution of LAI, plant coverage and the senescence in autumn;

COSMO model contains the phenology cycle is based on a 6-

2) year climatology. Cycle follows the same sinusoidal fitted curve between its max and min value each year while neglecting any influence, feedback on the environmental conditions;

Evolution of land surface modelling

1st generation
LSM:

2nd generation
LSM:

3rd generation
LSM:

Vegetation
dynamics

★ “bucket model”

★ “Big-leaf” approach

★ Photosynthesis

★ No explicit
treatment of
vegetation

★ Stomatal
conductance


★ Carbon cycle

(biogeography)


1970

1980

1990




Bucket model:
Manabe, 1969




BATS:
Dickinson, 1984

SIB:
Sellers et al., 1986
TERRA-ML



SiB2:
Sellers et. al, 1992

LSM:
Bonan, 1995



IBIS:
Foley et al., 1996

CLM

Seller et. al. (1997) classification

Vertical discretization

no canopy layer

“Big-leaf approach”

“2-leaf” approach

Multi-layer canopy

“bucket hydrology”

2-layer hydrology

Multi-layer hydrology


1970

1980


1990

2000


2010




Bucket model: Manabe, 1969



BATS:
Dickinson, 1984



CLM3:
Oleson et al, 1984



CLM 4.5:
Oleson et al, 2013

ORCHIDEE:

Ryder et al, 2013

8

Edouard Davin, 2013 – Introduction to the CLM

COSMO General Meeting, 2 - 7 September 2020

Parameterization of the surface fluxes

$$E_b + \sum_{k=1}^{ke_{soil,hy}} T_{rk} + E_i + E_{snow} = -(F_{qv}^3)_{sfc}$$


E_{snow} – evaporation from snow

E_b – evaporation from soil

T_{rk} – plant transpiration

E_i – interception

Biological processes play a major role in controlling evapotranspiration



2nd generation Biophysical models (*TERRA_ML*)

Stomatal behavior represented based on empirical relations (Jarvis et. al., 1976)




Figure: Bonan, 2002

3rd generation Photosynthesis models (*CLM*)

Stomatal conductance explicitly related to photosynthetic assimilation model using Ball-Berry approach (Collatz et. al., 1991)






Figure: Sellers et. al, 1997

Limitation of 2nd generation LSMs

Vegetation explicitly **represented** in 2nd generation LSMs **but...**

Stomatal conductance is **calculated empirically without** considering the actual process **controlling stomatal functioning**




Maximization of water use efficiency (photosynthesis/water)

Edouard Davin, 2013 – Introduction to the CLM

Stomatal functioning

Stomata open



High light levels

Moist leaf


Warm temperature

Moist air

Moderate CO₂

High leaf nitrogen

Stomata close (smaller pore opening)



Low light levels

Dry leaf

Cold temperature

Dry air

High CO₂

Low leaf nitrogen

Comparison

TERRA_ML

vs

New scheme

Surface temperature and energy balance

- Single interface with one temperature (***t_g***) and bulk density

- Distinguishes temperature and energy fluxes for canopy (***tv***) and ground (***tg***)

Radiation fluxes

- Fluxes based on grid scale albedo and temperature

- Albedo dependences from vegetation

- Technically in ***src_radiation***

Comparison

TERRA_ML

vs

New scheme

Stomatal conductance

BATS-based

Empirical Jarvis-type approach

Ball-Berry approach

Coupling with photosynthesis

“2-leaf” canopy
with diffuse/direct light

Subgrid-scale heterogeneity

Only accounts for partial
coverage of snow

Tile approach in ICON-TERRA

Tile approach

Vegetation types adapted for ICON

Individual Sub-Tasks 1

Implementation of three new modules in current
photosynthesis/phenology scheme of COSMO/ICON code

- 1) The canopy photosynthesis and stomatal regulation;
- 2) Carbon allocation and plant growth;
- 3) Autotrophic/Heterotrophic respiration and litter/soil carbon dynamics;

*The new modules will be implemented in a way to minimize the additional computational time introduced by these new parameterizations

Individual Sub-Tasks 2

Quantification of impact on implemented parameterizations

- * Impact of new scheme on the albedo and on evapotranspiration/latent heat flux will be quantified
- ** The start from a specific date need to be accounted for NWP applications, and the necessary predictions to be written out and read in each time step

Individual Sub-Tasks 3

Validation of implementation of new photosynthesis/phenology scheme

Sub-Task 3 includes verification of prognostic and diagnostic parameters affected by the photosynthesis/phenology scheme.

* Especially, the diagnostic parameters such as the 2m air temperature will be affected by an improved photosynthesis/phenology scheme.

** Adjustments to such diagnostics will be made if required.

Individual Sub-Tasks 4

Documentation of new photosynthesis/phenology scheme

In the final phase 4 required documentations in forms of internal works reports (COSMO consortium) as well as scientific publications for peer-reviewed journals will be prepared.

Our contacts:

GitHub page: <https://github.com/users/merajtoelle/projects/1>

Address: Center for Environmental Systems Research
Wilhelmshöher Allee 47, 34117 Kassel

Email: evgenychur@gmail.com

Background

- ➔ Vegetation dynamics are prescribed in lookup tables or in gridded maps with effective plant properties for each raster cell.
- ➔ As a consequence, the development of the leaf area index (LAI), for example, is the same in each year.
- ➔ The dynamics are “frozen” and hence do not depend on the prevailing weather conditions.
- ➔ Errors in turbulent fluxes resulting from this coarseness translate into changes in regional air and surface temperature, evolution of the atmospheric boundary layer, and spatial distribution of rainfall.
- ➔ In general, a warmer climate will accelerate vegetation development...


Differences between the models

	TERRA-ML	Veg3D	JS-BACH	LPJmL	Caraib
Vegetation layer	NO / Vegetation parameters	YES / Big leaf concept	YES / Big leaf concept	YES / Big leaf concept	YES / Big leaf concept (shaded and sunlit leaves)
Turbulent fluxes	Surface temperature	Vegetation temperature	Vegetation temperature	Vegetation temperature	Vegetation temperature
Vegetation parameters	Weighted average	Dominant			
Radiation	Albedo constant	Albedo depends on vegetation			

Modifications in TERRA

- modify...
- Add photosynthesis, respiration, allocation, and phenology

Gantt chart



FTEs summary

	Institution	CESR	MeteoSwiss	DWD
Task				
1		1.39	0.005	0.005
2		0.29	0.005	0.005
3		0.19	0.005	0.005
4		0.09	0.005	0.005
Total FTEs		1.96	0.02	0.02

*** Total of 2 FTEs over 2 years**