

# A new leaf phenology for the land surface scheme TERRA of the COSMO atmospheric model

**Jan-Peter Schulz<sup>1,3</sup>, Gerd Vogel<sup>2</sup> and Bodo Ahrens<sup>3</sup>**

<sup>1</sup>Biodiversity and Climate Research Centre (BiK-F), Frankfurt

<sup>2</sup>Deutscher Wetterdienst, Lindenberg

<sup>3</sup>Goethe Universität Frankfurt

COSMO / CLM / ART User Seminar, 17 - 19 Mar. 2014, Offenbach


# What is phenology?

Phenology is the study of periodic plant and animal life cycle events and how these are influenced by seasonal and inter-annual variations in climate, as well as habitat factors (such as elevation).

Wikipedia, 4 Mar. 2014

## Phenological stages in the production of almonds include

- winter chilling
- spring heat forcing flowering and leaf-out
- pollination of flowers
- development and growth of fruits during spring and summer



<http://treephenology.ucdavis.edu/>

# What is phenology?


Phenology is the study of periodic plant and animal life cycle events and how these are influenced by seasonal and inter-annual variations in climate, as well as habitat factors (such as elevation).

Wikipedia, 4 Mar. 2014

## Phenology depends on:

- Region


## Begin of flowering 2013



24.4. 27.4. 30.4. 3.5. 6.5. 9.5. 12.5. 15.5. 18.5.

Deutscher Wetterdienst (erstellt 09.01.2014 09:31 UTC)  
Kontakt: Landwirtschaft@dwd.de  
Geobasisdaten © Bundesamt für Kartographie und Geodäsie (www.bkg.bund.de)

## Fall of leaves 2013



15.10. 21.10. 27.10. 2.11. 8.11. 14.11. 20.11. 26.11.

**Apple  
tree**

The flowering of the apple trees starts in the southwest of Germany and progresses towards the northeast. The fall of leaves evolves roughly in opposite direction.

# What is phenology?

Phenology is the study of periodic plant and animal life cycle events and how these are influenced by seasonal and inter-annual variations in climate, as well as habitat factors (such as elevation).

Wikipedia, 4 Mar. 2014



## Phenology depends on:

- Region
- Year

# Begin of flowering

# Fall of leaves

## Deviation from multi-annual mean (in days)



**Apple tree**

Deutscher Wetterdienst (erstellt 09.01.2014 09:31 UTC)  
Kontakt: Landwirtschaft@dwd.de  
Geobasisdaten © Bundesamt für Kartographie und Geodäsie (www.bkg.bund.de)

There is a high inter-annual variability of the dates of flowering and fall of leaves of apple trees. In spring, there is a trend towards earlier dates of occurrence.

# Beginning of the phenological early spring

Period 1961-1990

Day of Year: 080



<http://www.dwd.de/>

© Geobasisdaten – Bundesamt für Naturschutz

The phenological early spring starts in the southwest of Germany and progresses towards the northeast. This is also the case for the whole of Europe.



# Beginning of the phenological early spring

Period 1961-1990

Day of Year: 090



Period 1980-2009

Day of Year: 90




<http://www.dwd.de/>

© Geobasisdaten – Bundesamt für Naturschutz

In recent years the phenological early spring shows a clear tendency to arrive earlier. This can be explained by the increase of spring temperatures.


## Inter-annual variability at Lindenberg




**Currently the seasonal cycle of leaf area index (LAI) is prescribed as climatology, based either on**

- a sinusoidal fit between a minimum and maximum value of LAI


## Leaf area index (grass)



## Leaf area index (grass)




## Leaf area index (grass)



## Currently the seasonal cycle of leaf area index (LAI) is prescribed as climatology, based either on

- a sinusoidal fit between a minimum and maximum value of LAI
- monthly mean values of LAI based on satellite retrievals





**Currently the seasonal cycle of leaf area index (LAI) is prescribed as climatology, based either on**

- a sinusoidal fit between a minimum and maximum value of LAI
- monthly mean values of LAI based on satellite retrievals

The model can **not account for inter-annual variations** of the growing season.


**Therefore ...**

... we need a **phenology scheme** in the COSMO / COSMO-CLM atmospheric model which allows the vegetation to adapt to the simulated seasonal and inter-annual variations in weather and climate, as well as to habitat factors (such as elevation).

# Phenology is governed, or limited, by:

- Temperature
- Day length
- Water availability
- NPP (net primary productivity), used e.g. in JSBACH, CLM, ORCHIDEE ...

# Governing phenology



denotes primary inputs  
denotes processes modeled  
denotes primary outputs  
 ----- Indicates linkage between models.

Dashed lines indicate linkages between the hydrological model and the vegetation module.  
 A – LAI affects transpiration from soil and groundwater stores.  
 B – LAI affects canopy interception, storage, and evaporation from canopy leaves.  
 C – Soil moisture affects photosynthesis.  
 D – The coupling between photosynthesis and stomatal conductance is used to estimate transpiration.

Arora (2002)

Example for the coupling between a hydrologic model and a dynamic vegetation module. The two primary variables exchanged between the two models are leaf area index and soil moisture.

# Phenology is governed, or limited, by:

- Temperature
- Day length
- Water availability
- ~~NPP (net primary productivity)~~

## Two approaches for phenology not depending on NPP adopted from:

- Polcher, J. (1994), *Thèse de doctorat, Univ. Pierre et Marie Curie, Paris*
- Knorr, W., et al. (2010), *J. Geophys. Res.*, **115**, G04017

## Phenology determining temperature

$$T(t) = \frac{\int_{-\infty}^0 T_S(t + \tilde{t}) e^{\tilde{t}/t} d\tilde{t}}{\int_{-\infty}^0 e^{\tilde{t}/t} d\tilde{t}}$$

This is equivalent to an exponentially declining memory of the plants for the surface temperature  $T_S$ .  $t$  is the averaging period for  $T_S$ .

# Phenology as function of temperature

## based on Polcher (1994)

$$\text{LAI}(t) = \begin{cases} \text{LAI}_{\min} & \text{if } T(t) \leq T_1 \\ \text{LAI}_{\min} + \frac{T(t) - T_1}{T_2 - T_1} (\text{LAI}_{\max} - \text{LAI}_{\min}) & \text{if } T_1 < T(t) \leq T_2 \\ \text{LAI}_{\max} & \text{if } T_2 < T(t) \end{cases}$$


$T_1$  : minimum limiting temperature

$T_2$  : maximum limiting temperature


$\text{LAI}_{\min}$  ,  $\text{LAI}_{\max}$  : minimum and maximum value of LAI



# Inter-annual variability at Lindenberg



## Leaf area index (grass)



based on Polcher  
(1994)



# Phenology as function of temperature

## based on Knorr et al. (2010)


$$\frac{dLAI(t)}{dt} = \begin{cases} k_{grow}(LAI_{max} - LAI(t)) & \text{if } T(t) \geq T_{on/off} \\ k_{shed}(LAI_{min} - LAI(t)) & \text{else} \end{cases}$$

$T_{on/off}$  : leaf onset and offset temperature

$k_{grow}$  ,  $k_{shed}$  : growth rate and shedding rate


$LAI_{max}$  ,  $LAI_{min}$  : maximum and minimum value of LAI

## Leaf area index (grass)






based on Knorr et al. (2010)

## Latent heat flux




## Latent heat flux







## Latent heat flux




## Latent heat flux



## Latent heat flux



## Latent heat flux



## Conclusions

- With the current parameterization TERRA can not account for the inter-annual variability of the phenology.
- Two approaches based on Polcher (1994) and Knorr et al. (2010) for simulating the seasonal cycle of phenology as function of temperature were implemented.
- The first one improves the simulations, the second one even gets very close to the observations of latent heat flux.
- The approach by Knorr et al. (2010) appears to be favourable due to the use of the concept of growth and shedding rates.
- The next steps are the extension of the scheme to more vegetation types, e.g. trees (deciduous and evergreen), and the implementation into the three-dimensional coupled model code.